Comparison of Actual Sickness with the Expectation according to the M.U. 1866-70 Experience, in Quinquennial Age-periods.

Age.	Years of		Actual Sickr	ness (Weeks).	Expected Sickness (Weeks).					
	Life at Risk.	First Six Months.	Second Six Months.	After Twelve Months.	Total.	First Six Months.	Second Six Months.	After Twelve Months.	Total.	
Under 25	11,817.5	7,963	477	851	9,291	8,034	397	228	8,659	
25-30	11,908.0	6,958	505	715	8,178	8,442	601	580	9,623	
30-35	11,632.5	7,577	622	1,221	9,420	9,122	866	961	10,949	
35-40	10,748.5	7,783	564	1,007	9.354	9,143	856	1,475	11,474	
40-45	8.434.5	7,091	835	2,755	10,681	8,496	993	1,779	11,268	
45-50	6,576.5	7,097	983	3,548	11,578	7,242	1,169	2,257	10,668	
50-55	4.912.0	6,620	1,274	4,358	12,252	6,765	1,317	2,642	10,724	
55-60	2.584.5	4,064	905	5,232	10,201	4,455	1,064	2,247	7,766	
60-65	1,325.5	2,652	596	4,113	7,361	2,955	906	2,122	5,983	
65-70	397.0	1,057	256	3,011	4,324	1,207	413	1,192	2,812	
70 and up-	1 1111	660	352	3,011	4,023	860	391	1,693	2,944	
wards				.,	,					
All ages	70,558.5	59,522	7.319	29,822	96,663	66,721	8.973	17,176	92,870	

Comparison of the Average Sickness per Member with the Expectation according to the M.U. 1866-70 Experience, in Quinquennial Age-periods.

		. !		New Zealan	d Experience.		English Experience.				
	Age.			Second Six Months.	After Twelve Months.	Total	First Six Months.	Second Six Months.	After Twelve Months.	Total.	
	•						1	.			
Under 25			0.68	0.04	0.07	0.79	0.68	0.03	0.02	0.73	
25-30			0.58	0.04	0.06	0.68	0.71	0.05	0.05	0.81	
30-35			0.65	0.05	0.11	0.81	0.78	0.08	0.08	0.94	
35-40			0.73	0.05	0.09	0.87	0.85	0.08	0.14	1.07	
40-45			0.84	0.10	0.33	1.27	1.01	0.12	0.21	1.34	
45-50			1.08	0.14	0.54	1.76	1.10	0.18	0.34	1.62	
50-55			1.35	0.26	0.88	2.49	1.38	0.26	0.54	2.18	
55-60			1.57	0.35	2.03	3.95	1.72	0.41	0.87	3.00	
60-65			2.00	0.45	3.10	5.55	2.23	0.68	1.60	4.51	
65-70			2.66	0.64	7.59	10.89	3.04	1.04	3.00	7.08	
70 and upv			2.97	1.58	13.57	18.12	3.87	1.76	7 63	13.26	

Comparison of Actual Mortality with the Expectation according to the M.U. 1866-70 Experience, in Quinquennial Age-periods.

					Number	of Deaths.	Death-rate per Thousand.		
	A	ge.	/ ÷ .		Actual.	Expected.	Actual.	Expected	
Under 25					56	74.04	4.74	6.27	
25-30					55	88.82	4.62	7.46	
30-35					55	97.58	4.73	8.39	
35-40	• •				53	108 02	4.93	10.05	
40-45					45	102.22	5.34	12.12	
45-50					51	96.94	7.75	14.74	
50-55					72	94.41	14.66	19.22	
55-60					54	66.87	20.89	25.87	
30-65					19	48.09	14.33	36.28	
6570					16	19.98	40.30	50.33	
70 and upwa	ards	••	• •	••	18	19.55	81.08	88.06	
A	ll ages				494	816.52	7.00	11.57	

Table of Secessions among those who were Sick and not Sick respectively, during the Quinquennium preceding Valuation, in Quinquennial Age-periods.

			<u>+</u>		Number	u of N	Iambara wh	<u> </u>					
Age at Beginning of Quinquennium or at Entry.		Number of Members who											
		A	dmitted before	ore Quinque	nnium.	Admitted during Quinquennium.							
			ek during ennium.	Were not Sick during Quinquennium.		ing nium.	Were Sick during Quinquennium.		Were not Sick during Quinquennium.		ing itum.		
		Lapsed during Quinquen- nium.	Were included in the Valuation.	Lapsed during Quinquen- nium.	Were included in the Valuation.	Died during Quinquennium.	Lapsed during Quinquen- nium.	Were included in the Valuation.	Lapsed during Quinquen- nium.	Were included in the Valuation.	Died during Quinquennium		
Under 25		161	751	811	634	39	121	972	1,437	3,001	32		
25-30		98	758	711	7 58	40	43	341	569	1,133	17		
30-35		98	857	586	833	44	30	191	311	689	11		
35-40		81	882	398	780	46	16	108	150	326	5		
40-45		43	780	229	583	49	5	43	34	124	2		
45-50	٠.	34	697	140	450	61	1	11	8	32	1		
50-55	• •	18	449	69	256	63				4	1		
55-60	• •	1	258	19	138	39			1	1			
60-65	••		118	7	59	20		1					
65–70	• •	••	32	2	9	8	••	• •	••	•••			
70 and upward	ls	••	20	••	4	16	••	••	••	•••			
All ages		534	5,602	2,972	4,504	425	216	1,667	2,510	5,310	69		