

Relation between Age and Frequency of each Group of Causes of Rejection for Military Service

This graph further illustrates the variation in frequency of each of the major groups of medical conditions which have caused men to be placed in Grade II or lower grades.

For example, disorders of the bones and organs of locomotion and cardio-vascular diseases are shown to increase rapidly with advance in age, whereas eye disorders do not show any marked variation over the range of ages covered by the graph.

This graph is based on Table 6 above, which gives similar information, though more finely classified as regards causes of rejection.




TABLE 7.—ANALYSIS OF HEIGHTS AND WEIGHTS OF VARIOUS CLASSES OF MEN MEDICALLY EXAMINED FOR MILITARY SERVICE

	Age, 18-19 Years.	Central Age, 28 Years.			Central Age, 42 Years.		
	Men placed in Grade I.	Men placed in Grade I.	Other Men.	Total.	Men placed in Grade I.	Other Men.	Total.
Height—							
Number of cases examined	1,915	510	216	726	312	1,149	1,461
Average measurement . .	5 8 $\frac{1}{4}$	5 8 $\frac{1}{4}$	5 7 $\frac{1}{4}$	5 7 $\frac{3}{4}$	5 7 $\frac{3}{4}$	5 7 $\frac{3}{4}$	5 7 $\frac{3}{4}$
Weight -							
Number of cases examined	373	Not available			154	448	602
Average measurement . .	10 1 $\frac{1}{4}$				11 2 $\frac{1}{2}$	10 12 $\frac{1}{2}$	11 1 $\frac{1}{2}$