Analysis of Deaths of Infants Under One Month, 1945
(Europeans only)

Cause of Death.	Under One Day.	One Day and Under One Week.	One Week and Under Two Weeks.	Two Weeks and Under Three Weeks.	Three Weeks and Under One Month.	Total.
Diphtheria		· · · · · · · · · · · · · · · · · · ·	 1	1 	1	2 2 1
Whooping-cough						
Influenza						
Syphilis					1	1
Convulsions						
Broncho-pneumonia		6	6	4	1	17
Pneumonia		3	1	1	1	6
Diarrhœa and enteritis		2		3	1	6
Congenital malforma-	21	49	22	7	4	103
tions				Ť	_	
Congenital debility	6	6	1			13
Injury at birth	33	71	11	4	i	$1\overline{20}$
Premature birth	244	131	16	4	2	397
Other diseases of early	36	57	10	$\overset{\bullet}{2}$	$\frac{z}{2}$	107
infancy	30	3,			_	101
Other causes	4	5	8	4	4	25
	- 4 .				-	
Totals—						
1946	344	331	76	30	18	799
1945	312	293	70	$\frac{30}{28}$	22	725
1944	289	$\frac{233}{270}$	76	43	14	692
1049	$\begin{array}{c} 253 \\ 254 \end{array}$	284	61	$\frac{49}{25}$	21	645
1040	260	269	54	$\frac{26}{26}$	$\frac{21}{25}$	629
1041	280	293	70	38	21	702
1941	200	±0.0	10	90	-1	102

Still-births.—The still-births for Europeans registered in 1947 numbered 911, giving a rate of 19·92 per 1,000 total births. This is the first occasion on which the still-birth rate has been below 20 per 1,000 births, the previous lowest figure (1946) being 21·75 per 1,000 total births. No figures are available for Maori still-births.

The following table shows the death-rates for still-births and deaths of infants under one month of age for the years 1942-47:—

Rates Per 1,000 Total Births

(Europeans only)

	Year.		Still-births.	Under One Day.	One Day and Under Two Days.	Two Days and Under One Week.	One Week and Under Two Weeks.	Two Weeks and Under One Month.
1942			25.85	$7 \cdot 54$	2.50	5 ·16	1.57	1.48
1943			$26 \cdot 25$	$8 \cdot 16$	$3 \cdot 53$	5 ·59	$1 \cdot 96$	$1 \cdot 48$
1944			$23 \cdot 23$	8.40	$2 \cdot 50$	5 ·35	$2 \cdot 21$	1.66
1945			$22 \cdot 84$	$8 \cdot 24$	$2 \cdot 43$	5.30	1.85	$1 \cdot 32$
1946			$21 \cdot 75$	8.04	$2 \cdot 90$	4.84	$1 \cdot 77$	$1 \cdot 12$
1947			$19 \cdot 92$	$7 \cdot 87$	2.89	4.44	1.31	1.20

Maternal Mortality.—Here again the Maori figures are not available.

For Europeans the maternal-mortality rate, including deaths from septic abortion, was 1.07 per 1,000 live births, as compared with 2.05 in 1946. After excluding deaths from septic abortion, the rate was 0.85 (1.76 in 1946).