Causes of Death-continued.

Class.	Causes of Death.	Males.	Females.	Total.	Class.	Causes of Death.	Males.	3	
	Orders and Diseases.					Orders and Diseases.		Females.	Total.
VI.—Local Diseases—continued.	Order 8.—Diseases of Reproductive System —continued (a.) Diseases of organs of generation—ctd. Diseases of uterus and vagina Disorders of menstrua- tion Pelvic abscess Perineal abscess		23 1 7	23 1 7 1	VII.—Violence.	Order 1.—Accident or Negligence. Fractures, contusions Gunshot wounds Cut, stab Burn, scald Sunstroke Poison Drowning Suffocation Otherwise	210 21 12 20 10 126 17 18	29 1 4 19 7 16 10 9	22 16 39 17 142
	Diseases of testes, penis, scrotum, &c (b.) Diseases of parturi-	•••	•••	•••		Total Order 1	425	95	520
	tion,— Abortion, miscarriage Puerperal mania Puerperal metritis Puerperal convulsions		20 1 	20 1 9		ORDER 2.—Homicide. Murder, manslaughter Wounds in battle Total Order 2	 3	₂	 5
	Placenta prævia (flood- ing) Phlegmasia dolens Other accidents of		17 1 31	17 1 31		ORDER 3.—Suicide. Gunshot wounds	24		24
	child-birth	1	118	119		Cut, stab Poison Drowning	15 5 7	2 8 7	17 13 14
	Total Order 8 ORDER 9.—Diseases of			 		Hanging Otherwise	13 5	$\begin{array}{c} 2 \\ 1 \end{array}$	15 6
	Organs of Locomotion. Caries, necrosis	3	1	4 4		Total Order 3	69	20	89
	Arthritis, ostitis Other diseases of organs of locomotion	3	3	6		ORDER 4.—Execution. Hanging	1	•••	1
	Total Order 9	9	5	14		Total Class VII	498	117	615
	Order 10.—Diseases of Integumentary System. Carbuncle Phlegmon, cellulitis Lupus	1 3	 2 2	1 5 2	NED AND NOT-CAUSES.	Dropsy Marasmus, &c Mortification, gangrene Tumour Abscess	162 1 3	126 1 1	288 1 4 1
	Ulcer, bed-sore Eczema Pemphigus Other diseases of inte-	 2 2 	₁	 3 2 	I.—Ill-defi Specified	Hæmorrhage Sudden (cause unascertained) Other ill-defined and	" 11 …	 2 	13
	gumentary system Total Order 10	8	5	13		not-specified causes Total Class VIII	177	130	307
	Total Class VI	_					4,689		

MORTALITY FROM FEBRILE AND ZYMOTIC DISEASES.

The deaths in 1905 from specific febrile or zymotic diseases amounted to 391, a proportion of 4.49 in every 10,000 persons living, and a decrease of 177 on the number of deaths in 1904, when the proportion was 6.72.

The feature of last year's mortality is the comparative freedom of the colony from some of the epidemic diseases most fatal to children—viz., measles, scarlet fever, scarlatina, and whooping-cough—the total deaths from these causes being only 21, against 58 in 1904 and 478 in 1903.