Causes of Death—continued.

Causes of Dealer Continuous									
	Causes of Death.		es.			Causes of Death.		les.	
Class.	Orders and Diseases.	Males.	Females.	Total.	Class.	Orders and Diseases.	Males.	Females.	Total.
VI.—Local Diseases—continued.	Order 8.—Diseases of Reproductive System —continued (a.) Diseases of organs of generation—ctd. Diseases of uterus and vagina Disorders of menstrua- tion Pelvic abscess Perineal abscess Diseases of testes, pe- nis, scrotum, &c	22	17  3 	17  5 	I.—Ill-defined and Not-specified Causes.	Order 1.—Accident or Negligence. Fractures, contusions Gunshot wounds Cut, stab Burn, scald Sunstroke Poison Drowning Suffocation Otherwise Total Order 1	210 24 12 28 1 8 161 33 29	25 1  19  4 25 20 5	12 47 1 12 186 53 34
	(b.) Diseases of parturition,— Abortion, miscarriage Puerperal mania Puerperal metritis Puerperal convulsions Placenta prævia (flood-	•••	13 1 1 13 15	13 1 1 13 15		Order 2.—Homicide. Murder, manslaughter Wounds in battle Total Order 2	4	2	6 
	ing) Phlegmasia dolens Other accidents of child-birth Total Order 8	 	44 ———————————————————————————————————	 44 ————————————————————————————————		ORDER 3.—Suicide. Gunshot wounds Cut, stab Poison Drowning	39 12 8 8	1 1 2 1	40 13 10 9
	ORDER 9.—Diseases of Organs of Locomotion. Caries, necrosis Arthritis, ostitis	5 2 2	7 1 3	12 3 5		Hanging Otherwise  Total Order 3  Order 4.—Execution.	19 6 92	3 2 10	
	Other diseases of organs of locomotion	9		20		Hanging Total Class VII	602	111	713
	Total Order 9  Order 10.—Diseases of Integumentary System. Carbuncle Phlegmon, cellulitis Lupus Ulcer, bed-sore Eczema Pemphigus Other diseases of integumentary system	1 7  2 2 1	2 7  2  1	3 14   4 2 2		Dropsy Marasmus, &c Mortification, gangrene Tumour Abscess Hæmorrhage Sudden (cause unascertained) Other ill-defined and not-specified causes	 202 1 3 2 1 11	 143   1 1	345 1 3 2 2 12
	Total Order 10 Total Class VI	l	12 سر	25		Total Class VIII	$\frac{220}{5,755}$	145	
	Total Class V1	2,771	2,142	+,913	<b>&gt;</b> (	General totals	o, 1004	E, 311	T0000

## MORTALITY FROM FEBRILE AND ZYMOTIC DISEASES.

The deaths in 1907 from specific febrile or zymotic diseases amounted to 1,185, a proportion of 12.89 in every 10,000 persons living, and an increase of 739 on the number of deaths in 1906, when the proportion was 4.98.

The diseases in this class that have caused the greatest mortality during the past ten years are stated in the table that follows. Of these, diarrheal complaints were by far the most fatal, causing 2,102 deaths in the decennial period. Influenza comes next, with 1,465 deaths; whooping-cough third, with 886 deaths; typhoid fever fourth, with 709 deaths; measles fifth, with 616