21

the population of those districts at the census of 1906, thus adding 22,133 to the foregoing figures, produces a total of 99,229. But, as it is very well known that several of the suburban districts have extended in a manner altogether beyond what can be accounted for by natural increase, the Town of Auckland has probably now considerably over one hundred thousand souls.

[Note.—The figures relating to Onehunga, though that borough is practically within a Greater

Auckland area, are not brought into the statistics which follow.]

Birth-rate.

The average birth-rates per 1,000 of the	populati	on for ter	n years 19	00-9 are,		
Auckland City						30.28
Auckland and suburban boroughs		• •	• •			28.28
The average death-rates per 1,000 of the	Death-		en vears l	900-9 ar	e. 	
Auckland City					٠,	12.97
Auckland and suburban boroughs				.: ••		11.28
Natural	Increase	of Popul	ation.		TO:	····· 1 000

Ditto per 1,000 of Mean Natural Births. Deaths. Increase Population. 795 484 18.48

1,279 Auckland City 839 233 606 Suburban boroughs 17.78Auckland and suburban boroughs 2,118 717 1,401 18.28

Both the death and birth rates are lowered in 1909; still, as compared with 1908, when the natural increase was 2.04 in excess of the previous year, 1909 shows an improvement. This improvement is only maintained by effecting reduction in the death-rate, and the due effect upon the population is almost neutralized by the continued fall in the birth-rate.

Infant Mortality.

In Auckland and suburban borough	s,—						
Deaths of infants under 1 year	were	• •			٠,	131	
Between 1 year and 5 years	• •		• • •			39	
						170	
T					ckland	Auckland and	
Deaths of children under 1 year per 100 births,—					City.	Suburban Boroughs.	
Mean for 5 years 1900-4					12.09	11.18	
Mean for 5 years 1905–9					9.34	8.37	
7/T . C . L			••,	• • •	10.71	9.77	
Mean for the whole D	ominio	10 for 10	years 189	9-1908	= 7.59.		

It will be noted from the above table that there is a satisfactory decrease in the infant-mortality rate in the Auckland District during the past ten years. The present year is the lowest in that period. In 1909 the deaths in the suburban boroughs were 35 only, and these are only 4.05 per 100 births

-a very low percentage.

Causes of Deaths.

Zumotic Diseases.

Deaths in Auckland	and su	burban bor	\mathbf{oughs}		• •		74	
For the last five years: 1905 1906 1907	•••	Deaths 45 42 103		1908 1909	• •	••	Deaths 139 74	3.
These 74 deaths were fr	om,							
Diarrhœal diseases							48	
Influenza	• •					• •	\dots 2	
Typhoid fever						• •	5	
${f Diphtheria}$		• •	• •	• •	• •		6	
${\bf Whooping\text{-}cough}$. • •	• •	• •		• •		5	
${\bf Croup} \qquad \dots$	• •	• •	• •	• •	• •	• •	\dots 2	
Other zymotic disea	ises	• •	• •		• •	• •	6	
			Cancer.				74	
Deaths in Auckland	l and su	burban bor	oughs	••	••		76	
For the last five years,-		Deaths.					Death	s.
1905		35		1908			55	
1906		49	i	1909	• •		76	
1907		50	- 1					