L	acerated P	erinæum.		Primip.	Multip.	Instrum.	
First degree					59	37	28
Second degree					19	6	. 2
Third degree					13	. 1	1
Other lacerations	••	• •	••	••]	••	• •	-

-			COMPLIC	CATIONS.
	No.	Alb. pres. on Dis- charge.	Alb. abs. on Dis- charge.	Accompanying Complications.
Albuminuria before labour	62	No. 11 (1 death)	No. 35	4 eclampsia; 2 nephritis; 1 mitral disease; 2 marginal placenta prævia; 1 hydramnios; 1 purulent eczema; 1 phlebitis; several retained membranes; 1 macerated fœtus; 3 with vision
With toxic symptoms	19	9	••	impaired, headache, vomiting, and loss of memory-cedema. 1 threatened puerperal mania; some with varicose veins; 2 stillbirths; several with anæmia.
Eclampsia		6	eighth hæmo covere	ellington admitted moribund; I labour induced at month. 2 at Auckland; I admitted for ante-partum rrhage, five hours after developed eclampsia, red after induced labour; I case of twins; I laboured at thirty-seventh week; 3 fits after delivery.
	<u> </u>		·	
	Cen- tral.	Marg. with Hæm.	Without Hæm.	Remarks.
Placenta prævia	No. 2	No. 9	No. 2	In two cases digital dilatation was used, both children alive; 3 children stillborn.

MORBIDITY LIST. (With Day of Occurrences of Temperature.)

m		Days.		
Temp. for at least Two Consecutive Days.		2nd. 3rd. 4th. 5th. 6th. 7th. 8th. 9th.	Some Causes.	
100 to 100·8	11	2 2 1 1 2 2	10 mastitis; 8 influenza; 11 sapræmia; 2 septicæmia; 5 scarlet fever; 1 purulent eczema; 3 constipation; 3 phlebitis.	
100·8 to 102	20		2 mania; 2 auto-infection; 1 mucus colitis; 1 pyelitis; 1 retained chorion blocking cervix.	
102 to 103	21	6 6 4	4 heavy albumen with anæmia; 4 adherent	
Over 103	16	1 5 4 2 3 3 3 4	placenta. 1 admitted fourth day with salpingitis.	