Cause.	· · · · · · · · · · · · · · · · · · ·	In Camp.	Out of Camp.	Total.	
	FI	EATHERSTON CA	AMP.	THE RESIDENCE OF THE PARTY OF T	
Influenza infection .	1	165	8	173	
Cerebro-spinal menin	igitis	4	2	6	
A 4.1			1 :	1	
Tubercle of lung			1	1	
		. 1		1	
Diabetes		• •	1	1	
Suicide		1		.]	
Accidental death		1	1	2	
Alcoholic poisoning		• •	1	1	
Totals		172	15	187	
		TRENTHAM CAM	MP.		
Influenza infection	!	73	14	87	
Cerebro-spinal menir	ngitis	$\cdot 2$		$rac{2}{2}$	
Valvular disease of h	neart	2	1	2	
Apople x y		3		3	
α .		1		1	
Accidental death		1		1	
Suicide		1	••	1	
Totals		83	14	97	
		AWAPUNI CAM	P.		
Influenza infection	[5	2	7	
tm 1 1 61		• •	1	1	
Totals		5	3	8	
	N	ARROW NECK C	AMP.		
Influenza infection		14	3 !	17	

Note.—In Featherston Camp 4 deaths from cerebro-spinal meningitis are included in the 165 deaths from influenza infection; in Trentham Camp 5 deaths from cerebro-spinal meningitis are included in the 73 deaths from influenza infection.

It will be seen that there were only 2 deaths by suicide, as compared with 8 in 1917, 3 in 1916, and 4 in 1915.

Sickness.

Owing to difference in classification of the returns, comparative figures for the incidence rate of illness in the four camps are not accurate. For the first two quarters of the year the admission-rate to hospital at Featherston was below that of 1917. In the third quarter it rose from 4.2 per thousand average strength to 14.4 per thousand. This was due to the first wave of influenza. In the fourth quarter it rose to 58.4 per thousand, a result of the second wave.

At Trentham the rate was slightly higher throughout the first two quarters than in 1917. In the third quarter it rose to 36.5 per thousand, due to a very widespread first wave of influence.

the third quarter it rose to 36.5 per thousand, due to a very widespread first wave of influenza. In the fourth quarter it rose to 70.5 per thousand, due to the November second wave of influenza.

At Awapuni and Narrow Neck the sickness-rate in the last two quarters is similarly increased.

Infectious Disease.

In the following table is shown the incidence of various forms of infectious disease among men attached to the four camps in 1918 as compared with 1917:-

Disease.	Featherston. Tren		tham. Awapuni.		Narrow Neck.		Totals.			
	1917.	1918.	1917.	1918.	1917.	1918.	1917.	1918.	1917.	1918.
Cerebro-spinal meningitis	10	44	3	11					13	55
Measles	28	140	26	118					54	258
Diphtheria	11	9	8	3	2				21	12
Enteric fever	1	1	1				1		3	1
Scarlet fever	2	6	2	1					4	7
Chicken-pox	2	4	3	1					5	5
Erysipelas	71	2	. 1	2					2	4
Simple pneumonia	7	; 6	4	3	2	5	1		14	14
Broncho-pneumonia		320		203	١	27				550
Influenza		4,202		3,024		226		432		7,884
Totals	62	4,734	48	3,366	4	258	2	432	116	8,790