

ABSTRACT OF TABLE 1, SHOWING THE ADMISSIONS INTO HOSPITALS, DEATHS, AND AVERAGE CONSTANTLY SICK AMONGST THE TROOPS IN THE VARIOUS CAMPS AND STATIONS FOR THE YEAR 1918.

Camp or Station.	Featherston.			Trentham.			Narrow Neck.			Awapuni.			Hanmer.			Rotorua.		
Average strength	5,674			2,744			266			396			29			72		
Average sick-time to each soldier (days)	9.56			21.97			23.16			7.46			24.16			8.62		
Average duration of each case of sickness (days)	9.56			12.22			12.49			8.21			12.51			9.69		
Diseases.	Admitted.	Died.	Average constantly Sick.	Admitted.	Died.	Average constantly Sick.	Admitted.	Died.	Average constantly Sick.	Admitted.	Died.	Average constantly Sick.	Admitted.	Died.	Average constantly Sick.			
<i>General Diseases.</i>																		
Group A—																		
Cerebro-spinal fever	36	10	2.60	6	6	0.47												
Chicken-pox	4	..	0.19	5	..	0.17												
Cow-pox	..	..	..	..	..	..				1								
Diphtheria	4	..	0.32	..	..	..												
Enteric fever	1	..	0.01	..	..	..												
German measles	55	..	1.20	65	..	2.40												
Measles	83	..	3.23	45	..	2.81												
Influenza	4,522	157	107.30	3,227	79	99.80				253	6	5.59		3	1.49			
Mumps	..	..	..	11	..	0.38				..	..	..		..	..			
Scarlet fever	5	..	0.38	1	..	0.08				..	..	..		..	..			
Whooping-cough	..	..	..	1	..	0.13				..	..	..		..	..			
Totals	4,710	167	115.23	3,361	85	106.24				254	6	5.61		43	3			
Group C—																		
Malaria	3	..	0.10	..	..	..				..	..	..		..	..			
Totals	3	..	0.10	..	..	..				..	..	..		..	..			
Group D—																		
Pyrexia of uncertain origin	4	..	0.23	2	..	0.04				11	..	0.34		1	0.03			
Totals	4	..	0.23	2	..	0.04				11	..	0.34		1	0.03			