The admissions numbered 1,109 (m. 553, f. 556), or 40 more than in the previous year. Of these, 161 had been previously under care, making the proportion of readmissions 14-52 per cent., and 948 patients were admitted for the first time.

The ratio to population of all admissions (exclusive of Maoris) was 7.07 (m. 7.34, f. 7.20) to 10,000, and of first admissions 6.17 (m. 6.19, f. 6.14), so that 1,388 persons in the general population contributed

one patient, and 1,620 contributed a patient admitted for the first time.

The discharges (excluding transfers) numbered 471, or 91 more than in 1935. 201 (or 74 more) harmless unrecovered persons were returned to the care of friends; and 270 (m. 121, f. 149) recovered —17 more than last year—representing a percentage of 24·34 (m. 21·88, f. 26·80) on the total admitted. With voluntary boarders added the percentage rises to 27·34. Altogether, 50·92 per cent. of the inmates admitted were able to leave institutional care.

Of a total of 8,796 patients under care, 465 (m. 252, f. 213) died, or 6.38 per cent. on the average number resident. An inquest is held in the case of every death, whatever the cause. The causes are detailed in Table VI, and the following is the percentage of causes mainly contributing: Senile decay, 18.71; diseases of the brain and nervous system group, 23.01; heart-disease, 19.57; tuberculosis, 6.45.

In Table VII the principal causes assigned for the mental breakdown in the admissions are stated; but as a matter of fact they are merely approximations, and these, with the small numbers with which we have to deal, show such divergencies from year to year that the proportion assigned to any one cause in any one year cannot be assumed to be our average incidence. Causation is always complex, and the most potent factor is the individual. Hereunder the assignment causations in the table referred to are grouped and shown in their relative proportions:—

					Males.	Females.	Total.
Heredity					8.68	$6 \cdot 83$	7.76
Congenital					$19 \cdot 89$	$12 \cdot 41$	$16 \cdot 14$
Predisposed by pr	evious at	$\operatorname{tack}$			$12 \cdot 12$	18.70	$15 \cdot 42$
Ct 1 . 1 1 1 1					$6 \cdot 87$	$14 \cdot 21$	10.55
Child-bearing (pue	erperal, n	on-septic	e, and lac	tation)		$1 \cdot 99$	0.99
Mental stress					$18 \cdot 44$	$20 \cdot 32$	$19 \cdot 39$
Toxic, including-	-		м.	F.			
Syphilis			$4 \cdot 88$	1.44)	0.05	2.00	c 10
Alcohol			$3 \cdot 61$	0.72	9.95	$3 \cdot 06$	$6 \cdot 49$
Traumatic					0.54		0.27
Disorder of nervou	s system,	includin	g— м.	F.			
Epilepsy	٠		$2 \cdot 89$	$2 \cdot 52$	$5 \cdot 06$	3.78	$4 \cdot 42$
Other bodily affec	tions				8.50	$6 \cdot 65$	$7 \cdot 57$
Senility	• •				$9 \cdot 95$	$12 \cdot 05$	11.00
					100.00	${100 \cdot 00}$	100.00

## VOLUNTARY BOARDERS.

Year.		First Admission.		Not First Admission.		Total Admissions.		Transferred to Register of Patients.			Died.			Discharged.			Remaining on 31st December.						
			м.	F.	т.	м.	F.	T.	M.	F.	т.	м.	F.	T.	м.	F.	т.	м.	F.	T,	м.	F.	т.
1912		٠.	6	17	23	0	0	0	6	17	23	1	3	4	0	0	0	4	3	7	1	11	12
1913		• •	18	19	37	0	$^{2}$	$^2$	18	21	39	2	5	7	0	1	1	12	12	24	5	14	19
1914			17	19	36	3	2	5	20	21	41	7	5	12	1	1	$^{2}$	11	15	26	6	14	20
1915			15	17	32	1	$^{2}$	3	16	19	35	3	4	7	0	1	1	8	14	22	11	14	25
1916			13	23	36	5	8	13	18	31	49	4	4	8	1	2	3	14	14	28	10	15	25
1917			14	21	35	1	13	14	15	34	49	6	6	12	0	1	1	10	23	33	9	19	28
1918			23	38	61	5	11	16	28	49	77	1	4	5	2	<b>2</b>	4	15	30	45	19	32	51
1919		• •	31	39	70	5	19	24	36	58	94	3	3	6	0	2	2	26	42	68	26	43	69
1920		• •	26	38	64	16	12	28	42	50	92	3	4	7	1	2	3	33	33	66	31	54	85
1921		٠.	39	39	78	11	15	26	50	54	104	2	3	5	5	1	6	38	4.1	79	36	63	99
1922	• •		47	38	85	10	16	26	57	54	111	6	5	11	4	4	8	40	48	88	43	60	
1923	• •	• •	44	50	94	15	21	36	59	71	130	3	6	9	3	3	6	47	47	94	49	75	124
1924			38	44	82	16	20	36	54		118	4	8	12	6	3	9	39	48	87	54	80	134
1925			64	59	123	15	34	49	79	93	172	10	15	25	6	3	9	60	63	123	57	92	149
1926		٠.	71	60	131	11	<b>2</b> 9	40	82	89	171	10	11	21	7	2	9	49	71	120	73		170
1927			70		140	25	32	57	95	102	197	8	6	14	5	12	17	75	58	133	80	123	203
1928		٠.	120		214	37	39	76	157	133	290	8	10	18	10	8	18	102	97	199	117	141	258
1929	• •	٠.	102		165	50	43	93	152	106	258	11	3	14	9	9	18	119		229		125	
1930	• •		152		236	41	35	76	193	119	312	15	9	24	8	6	14	130		209	170	150	320
1931			166		264	43	36	79	209	134	343	11	14	25	11	7	18	159		261	198	161	359
1932			150	85	235	50	<b>4</b> 0	90	200	125	325	24	32	56	14	5	19	190	108	298	170	141	311
1933	• •	٠.	112	109	221	53	32	85			306	11	14	25	5	4	9	121	107	228	198	157	355
1934			132	90	221	56	33	89		123	311	7	11	18	10	6	16	152	99	251	217	164	~
1935			130	87	217	40	55	95	170	142	312	18	20	38	19	10	29	151	116	267		160	
1936	• •	٠.	98	99	197	62	44	106	160	143	303	21	20	41	7	10	17	137	111	248	194	162	356

It will be noted that there were 303 voluntary admissions in the year under review, that 248 were discharged, and that 41 boarders, representing 13.53 per cent. of the admissions, had to be transferred to the register of patients (having shown mental disorder in degree sufficiently pronounced and sustained to render it improper for them to be continued as voluntary boarders), while the proportion of deaths on average number of resident was 4.72 per cent. The column devoted to "Not First Admissions" records readmissions on a return of illness in those previously treated to recovery as voluntary boarders. The average daily number of voluntary boarders in 1936 was 360.